
No.



HHIB UNIHED SHAYIBS OF AMIBRICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

Urito-Tny Aorth America, Inc.

There has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXAMIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE PHERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, BEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY SLARS FROM THE DATE OF THIS GRANE, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC PETENDSHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A-PUBLIC REPOSITORY AS PROVIDED BY LAW, THE USE TO EXCLUDE OTHERS FROM SELLING THE VARIETY OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE PURPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT DEBYTHE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

POTATO

'FL 2006'

In Testimonn Microst, I have hereunto set my hand and caused the seal of the Hunt Huristy Frotection Office to be affixed at the City of Washington, D.C. this fifth day of June, in the year two thousand and eight.

Atlast:

De-3-

Commissioner
Plant Variety Protection Office
Agricultural Marketing Service

demond To Schafe

Secretary of Agriculture

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY - PLANT VARIETY PROTECTION OFFICE

The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Peperwork Reduction Act (PRA) of 1995.

APPLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE

Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426).

(Instructions and information collection burden statement	on reverse)				
1. NAME OF OWNER			2. TEMPORARY DESIGNATI EXPERIMENTAL NAME	ON OR	3. VARIETY NAME
Frito-Lay North America, Inc.			1995 191.2		FL 2006
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, and Count	ry)		5. TELEPHONE (include area	code)	FOR OFFICIAL USE ONLY
		•	972-334-3822		PVPO NUMBER
7701 Legacy Drive Plano, Texas 75024		:	6. FAX (include area code) 972−334−5965		200200127
- Wes					FILING DATE
7: IF THE OWNER NAMED IS NOT A "PERSON", GIVE FORM OF ORGANIZATION (corporation, partnership, association, etc.) Corporation	8. IF INCORPORAT STATE OF INCO Delaware		9. DATE OF INCORPORATION 8/8/89	N	03/29/02
10. NAME AND ADDRESS OF OWNER REPRESENTATIVE(S) TO SERVE IN THE	S APPLICATION. (First	t person listed will re	celve ell papers)		FILING AND EXAMINATION FEES:
Robert J. Jondle, Esq.					
Rothwell, Figg, Ernst & Manbeck 1425 K Street NW					, 2705.00
Suite 800					R DATE 0 3/29/02
Washington, D.C. 20005					CERTIFICATION FEE:
					\$ 768.00
•				<u>'</u>	DATE 09/07/07
11. TELEPHONE (Include area code) 12. FAX (Include area code)	13. E-M/	All		14 CROE	KIND (Common Name)
402-333-1550 402-333-1510		5 to 10 to 1	hwellfigg.com		otato
15. GENUS AND SPECIES NAME OF CROP	16. FAA	AILY NAME (Botanic	el)	17. IS TH HYBR	E VARIETY A FIRST GENERATION ID?
Solanum tuberosum	So	lanaceae]	☐ YES 🏅 NO
18. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow reverse) a. Exhibit A. Origin and Breeding History of the Variety	instructions on	CERTIFIED	OWNER SPECIFY THAT SEED C SEED? See Section 83(a) of VES (If "yes", answer items 20 and 21 below)	F THIS VAF the Plant Va	RIETY BE SOLD AS A CLASS OF unitary Protection Act) NO (If "no", go to item 22)
b. Ki Exhibit B. Statement of Distinctness c. Ki Exhibit C. Objective Description of Variety	•	20. DOES THE C	OWNER SPECIFY THAT SEED O	OF THIS	☐ YES ☐ NO
d.		VARIETY BE	LIMITED AS TO NUMBER OF C CH CLASSES?	LASSES7	<u> </u>
e. Exhibit E. Statement of the Basis of the Owner's Ownership f. Voucher Sample (2,500 viable untreated seeds or, for tuber propagat	ad variation		В точно] [] OCKITAED
verification that tissue culture will be deposited and maintained in an a repository)		21. DOES THE C VARIETY BE	WNER SPECIFY THAT SEED O	F THIS ENERATIO	NS? NO
g.	the United	IF YES, SPEC NUMBER 1,2	1 1	السا	REGISTERED CERTIFIED
22. HAS THE VARIETY (INCLUDING ANY HARVESTED MATERIAL) OR A HYBRIC	PRODUCED				
FROM THIS VARIETY BEEN SOLD, DISPOSED OF, TRANSFERRED, OR USE OTHER COUNTRIES?	ED IN THE U.S. OR	PROPERTY	RIGHT (PLANT BREEDER'S RIG	HT OR PAT	TY PROTECTED BY INTELLECTUAL FENT)?
YES 2-22-07 8-6 NO IF YES, YOU MUST PROVIDE THE DATE OF FIRST SALE, DISPOSITION, TF FOR EACH COUNTRY AND THE CIRCUMSTANCES. (Please use space indic	RANSFER, OR USE caled on reverse.)		(ES SE GIVE COUNTRY, DATE OF NUMBER. (Please use space in	_	NO ISSUANCE AND ASSIGNED REVERSE.)
24. The owners declare that a viable sample of basic seed of the variety will be furni	shed with application a	and will be replenishe	d upon request in accordance w		
for a tuber propagated vanery a tissue culture will be deposited in a public reposi- The undersigned owner(s) is(are) the owner of this sexually reproduced or tuber	silory and maintained fo occopadated plant vada	or the duration of the	certificate.		
and is entitled to protection under the provisions of Section 42 of the Plant Varie Owner(s) is(are) informed that false representation herein can jeopardize protect	ty Protection Act.		·		
SIGNATURE OF OWNER		SIGNATURE OF	OWNER	**************	4
Thomas P. Solur		-	•		•
NAME (Please print or type)		NAME (Please pri	int or type)		
Thomas P. Schur	. '			.*	
CAPACITY OR TITLE DATE		CAPACITY OR TI	TLE		DATE
Secretary, Recot, Inc.	SG SA				
S&T-470 (04-01) designed by the Plant Variety Protection Office with WordPerfect 6.0a	. Replaces STD-470 (02-99) which is obsc	Hete. (See reverse for inst	ructions and	i information collection burden statement)

INSTRUCTIONS

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed Exhibits A, B, C, E; (3) at least 2,500 viable untreater seeds, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in a public repository prior to issuance of a certificate; (4) check drawn on a U.S. bank for \$2,450 (\$300 filing fee and \$2,150 examination fee), payable to "Treasurer of the United States" (See Section 97.175 of the Regulation: and Rules of Practice.) Partial applications will be held in the PVPO for not more than 30 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 500, NAL Building, 1030: Baltimore Blvd., Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are sel explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$300 for issuance of the Certificate.

Plant Variety Protection Office Telephone: (301) 504-5518

#200200127

ITEM

16a. Give:

- the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified.
- 1.6b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences;
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 16c. Exhibit C forms are available from the PVPO for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 16e. Section 52(4) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. The applicant may be the actual breeder, the employee of the breeder, the owner through purchase or inheritance, etc.
- 17. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant may NOT reverse this affirmative decision after the variety has been sold and so labelled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See P.L. 103-349 for additional information.)
- 20. See Sections 41, 42 and 43 of the Act and Section 97.175 of the regulations for eligibility requirements.

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing a change of ownership or assignment is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131) 97.175(h) of Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant should check the variety names proposed by contacting: Seed Branch, AMS, USBA, Room \$\mathbb{P}\$13, Building 306, Beltsville Agricultural Research Center—East, Beltsville, MD 20705. Telephone: (301) 504-8089.

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gethering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Officer, OIRM, AG Box 7630, Washington, DC 20250; and to the Office of Management and Budget, Paperwork Reduction Project (OMB No. 0681-0066), Washington, DC 20503.

SD-470 (04-95) (REVERSE)

Date and Place of First Sale of FL 2006

'FL 2006' was first sold in South Africa in January 2005.

Exhibit A: Origin and Breeding History of the Variety

FL 2006 originated in the Frito-Lay, Inc. private potato breeding program. The variety is the result of classical hybridization breeding. No gene insertion was involved in the breeding of FL 2006 or its parents. In 1991, Robert W. Hoopes made a cross at the Frito-Lay Research facility near Rhinelander, Wisconsin between the varieties Atlantic and FL 1815. Atlantic was chosen as a breeding parent because of its high solids and yields and its potential for transmitting Golden Nematode resistance to its progeny. FL 1815 was chosen as a breeding parent because of its high yields, excellent chip quality out of cold storage and its potential for transmitting scab resistance.

Seeds from the cross Atlantic x FL 1815 were sown in a greenhouse near Rhinelander in the summer of 1994 and the resulting tubers were harvested in late fall of that year. Seedling tubers were planted in the field in the spring of 1995. One of the selections from this progeny was given the designation "1995 191.2". This selection was tested for two years in the fields near Rhinelander and one more year in Rhinelander, Iola and Florida, with solids measurements and potato chip fry samples taken after each harvest. It had excellent yields in the central Wisconsin trial out of the mild and late harvests. 1995 191.2 was found to have oval blocky yellow-fleshed tubers that flatten somewhat with size. Its outstanding attributes were excellent chip appearance when processed fresh from the field and over a seven month period from storage.

In 1998, 1995 191.2 was given the name "FL 2006". It was tested in twenty-six variety trials conducted by Frito-Lay throughout the United States, Canada and Europe in 1999 and 2000. In long growing season areas, it consistently had higher solids and yields than Snowden. It also continues to fry well out of extended storage. FL 2006 has completed two consumer tests at parity with Frito-Lay Gold Standard potato chips. Its average total defects have been less than 6, indicating excellent chipping quality.

FL 2006 was observed for 6 generations in 26 locations and was determined to be genetically uniform and stable from generation to generation with no evidence of variants.

Tissue culture plantlets of FL 2006 were established in 1998 and are maintained at the Frito-Lay Research facility near Rhinelander, Wisconsin with a duplicate maintained by the Wisconsin Seed Potato Certification Program in Madison, Wisconsin.






Exhibit B: Statement of Distinctness


As a chipping variety for fresh use and for up to seven months use out of storage at 52°F, FL 2006 is most similar to Snowden. FL 2006 can be distinguished from Snowden with regard to the following traits: FL 2006 tubers have pale yellow flesh (RHS 162C), while Snowden tubers have white flesh (RHS 158A). FL 2006 has tubers with violet eyes (RHS 87A), while the eyes of the tubers of Snowden do not contain pigmentation. FL 2006 has an average of 7.2 secondary and tertiary leaflet pairs, while Snowden has an average of 4 secondary and tertiary leaflet pairs. While both FL 2006 and Snowden have round tuber shapes, the tubers of FL 2006 tend to be more narrow and longer, while the tubers of Snowden are more blocky and frequently compressed. Additionally, the flower color of FL 2006 is blue-violet (RHS 96B fading to RHS 97B), while the flowers of Snowden are white (RHS 155C).

The isozyme pattern of FL 2006, as established by Dr. David Douches of Michigan State University is unique among known North American varieties. This is detailed in Exhibit D, Additional Description of the Variety.









Tuber flesh color RHS #158A, white

Tuber flesh color RHS # 162C, pale yellow

Tuber secondary skin color in eyes RHS #


87A, violet

Tuber secondary skin color in eyes absent, white


No. R590-15 WESTCOTTO Flexible




Corolla color RHS #155C, white




Corolla color RHS #96B when first open, fades to 97B



#200200127



#200200127



FL 2006

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY DIVISION PLANT VARIETY PROTECTION OFFICE

EXHIBIT C OBJECTIVE DESCRIPTION OF VARIETY POTATO (Solanum tuberosum L.)

Public reporting burden for this collection of information is estimated to average __minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the form. Send comments regarding this burden estimate or any other aspects of this collection of information, including suggestions for reducing this burden, to USDA, OIRM, Clearance Officer, AG Box 7630, Washington, DC 20259, regarding OMB No. 0581-0055. When replying, refer to OMB number and form number you your letter.

INSTRUCTIONS

#200200127

The Objective Description Form:

The objective description form lists characteristics to be used as the basis for developing the description of potato varieties. It is designed to guide the applicant in describing a variety in detail so a meaningful comparison with other potato varieties can be accomplished. It is recommended that this form be completed in as much detail as possible to ensure an accurate description. Please fill in the requested data and place the appropriate number that describes the varietal characters typical of this potato variety and the reference varieties in the respective boxes.

Test Guidelines:

Any statistical and trial (field test) data that may be necessary to support the variety description should be attached to this form. Please include for trial data the plot size, number of replications, number of plants, plant spacing, trial locations and growing periods. Trials should normally be conducted at one place, in the region that the variety has been adapted for, with a minimum of one growing period in the United States. All comparative data should be determined from varieties entered in the same trials. The size of the plots should be such that plants or parts of plants may be removed for measuring and counting without prejudice to the observations which must be made at the end of the growing period. As a minimum, each test should include a total of 60 plants which should be divided between two or more replicates. Separate plots for observation and measuring can only be used if they have been subject to similar environmental conditions. To determine color for a plant or plant parts a recognized standard color chart must be used such as the Royal Horticultural Society (R.H.S.) Color Chart.

Reference Varieties:

The application variety should be compared to at least one reference variety preferably a set of reference varieties. The reference varieties should be market class standard varieties currently grown in the United States and or the variety(ies) most similar. The following varieties are recommended as market class standards to be used as reference varieties:

Yellow-flesh table-stock	Yukon Gold
Round-white table-stock	Superior
Chip-processing	Atlantic, Snowden, Norchip
Frozen-processing	Russet Burbank
Russet table-stock	Russet Burbank, Russet Norkotah, Goldrush
	Red Pontiac, Red Norland, Red Lasoda

If the applicant does not use one of the recommended reference varieties the PVP office may not have a complete description for the reference variety used; therefore the applicant may have to supply this description by completing an Exhibit C form for the reference variety.

Characteristics:

The plant type and growth habit characteristics are collected at early first bloom. Figure 1 is supplied to help visualize the growth habit. For this descriptor, look at the stems rather than the stems and foliage. Plant maturity is measured at natural vine senescence.

Stem characteristics are also collected at early bloom. Stem anthocyanin coloration is divided into two descriptors: Location and intensity. Figure 12 is supplied to give an example of stem wings.

STD-476 (01-96)

Page 1 of 19

Leaf characteristics are observed at early first bloom. Fully-developed leaves located on the middle third of the plant should be used. Leaf pubescence refers to general trichomes. Figure 2 is supplied for examples of leaf silhouette. Figure 3 should be used to describe terminal and primary leaflet shape. Figures 4 and 5 are used to describe the terminal and primary leaflet shape of tip and base, respectively. To measure the total number of primary leaflets pairs, collect 10 fully- developed petioles (with leaves attached from each replication and take the average number of secondary and tertiary leaflets. Figure 11 is supplied to define leaf characteristics. Glandular trichomes should be described through descriptor #12 (Additional Comments and Characteristics). Leaf stipules are shown in figure 13 for visual definition.

Inflorescence characteristics should be measured at early first bloom. Figures 6 and 7 are supplied to describe corolla and anther shape, respectively. Corolla, calyx, anther, stigma and pollen should be observed on newly opened flowers.

Berry production should be based on field-grown plants rather than greenhouse plants.

Tuber characteristics should be observed following harvest. Figures 9 and 10 are available to describe distribution of secondary color and tuber shape, respectively.

Disease and pest reactions should be based upon specific tests rather than field observations. Other diseases or pests reactions not requested can be described if it is felt that it would be helpful to the description.

Quality characteristics should be described according to the market use.

If the plant is transgenic, this gene insertion(s) should be described.

Chemical identification and any other characteristics can be describe if they are helpful in distinguishing the variety.

A rating system of 1-9 provides a scale for describing most characteristics in this form. Characteristic may be rated with intermediate values where the characteristic grades gradually from one extreme to another. For example, if the character states are described as: 3 = Small; 5 = Medium; 7 = Large; the other values of 1, 2, 4, 6, 8, or 9 may be selected.

Legend:

V = Application Variety

R1-R4 = Reference Varieties

* = Both the reference variety(ies) and application variety must be described for characteristics designated with an asterisk.

AME OF APPLICANT(S)	we are a second or		FOR C	FFICIAL USE ONL	Y
Frito-Lay North Am	erica, Inc.		PVPO _{ANEN}	Т В 2 0 0 1	2 7
DDRESS (Street and No. or 1	R.F.D. No., City, State, and Z	ip Code)	VARIETY	(V) NAME	
+5000 Hopperd De So 7701 Legacy Drive			FL 2006		
Plano, Texas 75024	458		TEMPORA DESIGNAT	RY OR EXPERIMENTON	TAL
			1995 19	1.2	
REFERENCE VAR	RIETIES: Enter the	reference variety na	ıme in the	appropriate bo	×
Reference Variety 1 (R1)	Reference Variety 2 (R2	Reference Variety	3 (R3)	Reference Variety 4 (F	R4)
Snowden					
I. MARKET CHARACTERIS	TICS:				
MARKET CLASS:					
	stock; 2 = Round-white tabl 6 = Other	estock; 3 = Chip-processir	ng; 4 = Frozen	n-processing;	
v 3	R1 3 R	2 R3		R4	
2. PLANT CHARACTERIST	ICS:				
GROWTH HABIT: (See $3 = \text{Erect} (> 45^{\circ} \text{ with})$	figure 1) ground); 5 = Semi-erect (30	0-45° with ground); 7 = Sp	reading.		
v 5	R1 5 R	2 R3		R4	
TYPE:			· · · · <u>· · · · · · · · · · · · · · · </u>		
1 = Stem (foliage oper	n, stems clearly visible); 2 =	Intermediate; 3 = Leaf (F	Toliage closed,	stems hardly visible)	
v 2	R1 2 R 2	2 R3		R4	
			······································		
MATURITY: Days after turity of FL 2006 is	planting (DAP) at vine senes comparable to Snowd	cence en. Time to matur	rity varie	s according to	spec
	120			R4	
v 130	R1 130 R2	R3		<u> </u>	
V 130 PLANTING DATE:	R1 130 R2	R3	l	ality and grow	ing s
	R1 130 R2	R3	l	ality and grow	ing s
PLANTING DATE: V R1			l	11 1	ing s
PLANTING DATE:			loca	11 1	ing s

The

STD-476 (01-96)

Page 3 of 19

V 4	R1 4	R2	R3	R4
STEM CHARACTERIS	STICS: Measure at early f	irst bloom		
* STEM ANTHOCYAN	IN COLORATION:			
1 = Absent; 3 = V	Weak; 5 = Medium; 7 =	Strong; 9 = Very S	Strong	
V 7	R1 1	R2	R3	R4
STEM WINGS: (See f	igure 12)			
	Veak; $5 = Medium; 7 =$	Strong; $9 = \text{Very S}$	trong	
V 7	R1 7	R2	R3	R4
LEAF CHARACTERIST	TCS:			
LEAF COLOR: (Obse	rve fully developed leaves l	ocated on middle 1/3 oj	f plant)	·
1 = Yellowish-green	1; 2 = Olive-green; 3 = 1	Medium green; 4 =	Dark green; 5 = Grey-gr	cen; 6 = Other
v 4	R1 3	R2	R3	R4
,				
LEAF COLOR CHART	Γ VALUE: Royal Horticu	lture Society Color C	Chart or Munsell Color Ch	art
LEAF COLOR CHAR? (Observe fully developed	T VALUE: Royal Horticu	ulture Society Color (Chart or Munsell Color Chappropriate color chart)	art
(Observe fully developed V 147A	F VALUE: Royal Horticule leaves located on middle 1/	alture Society Color C 3 of plant & circle the	Chart or Munsell Color Chappropriate color chart)	R4
(Observe fully developed V 147A Dark	R1 137A	3 of plant & circle the	appropriate color chart)	
(Observe fully developed V 147A Dark LEAF PUBESCENCE I	R1 137A	3 of plant & circle the	appropriate color chart)	
(Observe fully developed V 147A Dark LEAF PUBESCENCE I	R1 137A DENSITY: parse; 3 = Medium; 4 =	R2 Thick; 5 = Heavy	appropriate color chart) R3	R4
(Observe fully developed V 147A Dark LEAF PUBESCENCE I 1 = Absent; 2 = Sp	R1 137A DENSITY: Parse; 3 = Medium; 4 =	3 of plant & circle the	appropriate color chart)	
(Observe fully developed V 147A Dark LEAF PUBESCENCE I 1 = Absent; 2 = Sp V 4 on back, 3 on LEAF PUBESCENCE I	R1 137A DENSITY: parse; 3 = Medium; 4 = R1 3 front LENGTH:	R2 Thick; 5 = Heavy	appropriate color chart) R3	R4
(Observe fully developed V 147A Dark LEAF PUBESCENCE I 1 = Absent; 2 = Sp V 4 on back, 3 on LEAF PUBESCENCE I	R1 137A DENSITY: parse; 3 = Medium; 4 = R1 3 front	R2 Thick; 5 = Heavy	appropriate color chart) R3	R4
(Observe fully developed V 147A Dark LEAF PUBESCENCE I 1 = Absent; 2 = Sp V 4 4 on back, 3 on LEAF PUBESCENCE I	R1 137A DENSITY: parse; 3 = Medium; 4 = R1 3 front LENGTH:	R2 Thick; 5 = Heavy	appropriate color chart) R3	R4
(Observe fully developed V 147A Dark LEAF PUBESCENCE I 1 = Absent; 2 = Sp V 4 On back, 3 on LEAF PUBESCENCE I 1 = None; 2 = Shor V 3	R1 137A DENSITY: Parse; 3 = Medium; 4 = R1 3 front LENGTH: t; 3 = Medium; 4 = Lo	R2 Thick; 5 = Heavy R2 ng; 5 = Very long R2	R3 R3	R4 R4
(Observe fully developed V 147A Dark LEAF PUBESCENCE I 1 = Absent; 2 = Sp V 4 On back, 3 on LEAF PUBESCENCE I 1 = None; 2 = Shor V 3	R1 137A DENSITY: Parse; 3 = Medium; 4 = R1 3 front LENGTH: t; 3 = Medium; 4 = Lo	R2 Thick; 5 = Heavy R2 ng; 5 = Very long R2	R3 R3	R4 R4
(Observe fully developed V 147A Dark LEAF PUBESCENCE I 1 = Absent; 2 = Sp V 4 On back, 3 on LEAF PUBESCENCE I 1 = None; 2 = Shor V 3 (Note: Descriptor #19)	R1 137A DENSITY: DATE: DENSITY: DATE: DENSITY: DATE: DENSITY: DATE: DENSITY: DENSIT	R2 Thick; 5 = Heavy R2 ng; 5 = Very long R2	R3 R3	R4 R4
(Observe fully developed V 147A Dark LEAF PUBESCENCE I 1 = Absent; 2 = Sp V 4 On back, 3 on LEAF PUBESCENCE I 1 = None; 2 = Shor V 3 (Note: Descriptor #19)	R1 137A DENSITY: Parse; 3 = Medium; 4 = R1 3 front LENGTH: Tt; 3 = Medium; 4 = Lo R1 4 Can be used to describe the (See figure 2)	R2 Thick; 5 = Heavy R2 ng; 5 = Very long R2	R3 R3	R4 R4
(Observe fully developed V 147A Dark LEAF PUBESCENCE I 1 = Absent; 2 = Sp V 4 4 on back, 3 on LEAF PUBESCENCE I 1 = None; 2 = Shor V 3 (Note: Descriptor #19)	R1 137A DENSITY: Parse; 3 = Medium; 4 = R1 3 front LENGTH: Tt; 3 = Medium; 4 = Lo R1 4 Can be used to describe the (See figure 2)	R2 Thick; 5 = Heavy R2 ng; 5 = Very long R2	R3 R3	R4 R4

MATURITY CLASS:

	Weak; 5 = Medium; 7		rong	
v 7	R1 1	R2	R3	R4
LEAF STIPULES SIZ 1 = Absent; 3 =	ZE: (See figure 13) Small; 5 = Medium; 7	= Large		
v 5	R1 7	R2	R3	R4
1 = Narrowly ovat	ET SHAPE: (See figure 3 te; 2 = Medium ovate; 5 Oblong; 8 = Other		Lanceolate; 5 = Ellipt	ical;
v 2	R1 2	R2	R3	R4
	TTIP SHAPE: (See figuuspidate; 3 = Acuminate		ner	
v 3	R1 3	R2	R3	R4
	T BASE SHAPE: (See fit Acute; 3 = Obtuse; 4 =		re; 6 = Lobed; 7 = Of	R4
	T MARGIN WAVINESS light; 3 = Weak; 4 = M			
The leaves tend to	R1 4/5 be quite textured	R2	R3	R4
NUMBER OF PRIMA	RY LEAFLET PAIRS: ((See figure 11)		
AVERAGE: V 4	R1 4	R2	R3	R4
RANGE: V 4 to 5 R1	4 to 5	to	R3 to	R4 to
	TIP SHAPE: (See figure spidate; 3 = Acuminate;		er	
v 3	R1 3	R2	R3	R4
D-476 (01-96)				Page 5 of 19

*	PRIMA	RY LEAI			- Mod	: <i>1</i>	- T	; 5 = Vei	_				***************************************		
. [o 4		3man, 3	3		R2	; 5 = Vei	- 7	ge R3			R4		
P	1 = 1	Varrowly	ovate;	APE: <i>(Sei</i> 2 = Medi long; 8 =	ium ovat	e; 3 =	Broadly	ovate; 4	= Lan	ceolate	; 5 = E	lliptica	 l;		
	v	1		R1	2		R2		÷	R3			R	4	
P	RIMARY 1 = C	LEAFL uneate;	ET BAS 2 = Acı	E SHAPI ite; 3 =	E: <i>(See j</i> Obtuse;	igure 5 4 = C	& 11) Cordate;	5 = Trunc	ate; (i = Lo	bed; 7 :	= Othe	r		*.
	v	4		R1	4		R2		{	R3			R	4	
N	UMBER AVERA		ONDAR	Y AND T	ERTIAI	RY LE	AFLET P	AIRS: (Se	e figu	re 11)		i		· · · · · ·	<u> </u>
	-	7.2		R1	4	[R2			R3			R	•	
RAN	_					7[1	f			<u></u>		
V	4 to	9	R1	3 to	5	R2	1	to	R3		to		R4	to)
5. IN	FLORES	CENCE	CHARA	CTERIS	TICS:										
N	MBER (OF INFL	ORESC	ENCE / I	LANT:										
	AVERA		Г			Г			۲		· · · · · · · · · · · · · · · · · · ·	٦			_,
Ĺ	V	4.8	L	R1	2	Ĺ	R2			R3]	R4		
RANG	GE:					, r			· —						
7 2	2 to	9	R1	1 to	2	R2	t	0	R3		to]	R4	to	
						<u></u>							"		····
			ETS / D	NFLORE	SCENC	E:									
ſ	AVERAG V	3E:	<u></u>	D1 //	.9	Γ	700		Γ	D 2					7
L	V	10	L	R1 4	.9	_	R2		L	R3			R4		
RAN	GE:		Г	·					1						
1	.0 to	34	R1	1 to	10	R2	to	<u> </u>	R3		to	[R4	to	
								yal Hortic			y Color	Chart (or Mun	sell Colo	r Char
v	96E		R1	l		R2			R3				R4		
		.nt-dar	k pig	ment											
∪-47 €	(01-96)													Page 6	of 19

*			R SURFACE COLOR CH		Royal Horticulture Society C	color Chart or Munsell Co	lor Chart
	\mathbf{v}	96C	R1 155C	R2	R3	R4	
STD	-476 (01-96)				Page 6 F	-5 40

TD-476 (01-96) Page 6.5 of 19

#200200127

1			ı	ı	l l				
v	3		R1 1		R2		R3		R4
	COROLLA 1 = Ve	A SHAPI ry rotate	E: (See figur ; 2 = Rotat	e 6) e; 3 =]	Pentagonal; 4 =	Semi-ste	llate; 5 = St	ellate	
v	3		R1 4		R2		R3		R4
			YANIN COI = Weak; 5 =		ION: .m; 7 = Strong;	9 = Ve	ry strong		
V	7		R1 1		R2		R3		R4
	ANTHER (COLOR (CHART VA opened flow	LUE: Fer is full	Royal Horticultur	e Society ircle the c	Color Chart	or Munse	ll Color Chart
V	12A		R1 14	В	R2		R3		R4
v	$1 = \mathbf{Bro}$	ad cone;	(See figure 7 2 = Narro R1 3 ter/even	w cone;	3 = Pear shape R2 anthers	cone; 4	= Loose; 5 :	= Other_	R4
V gma	1 = Broad 2 a little	shor	R1 3 ter/even	with	R2 anthers	cone; 4		= Other_	R4
V gma	1 = Broad 2 a little	shor	R1 3 ter/even	with	R2 anthers	cone; 4		= Other_	R4
V gma 1	2 a little POLLEN P 1 = Non-	short RODUCT e; 3 = S	R1 3 ter/even FION: ome; 5 = A	w cone;	R2 anthers	cone; 4	R3	= Other_	
V gma 1	2 a little POLLEN P 1 = Non-	RODUCTE; 3 = S	2 = Narro R1 3 ter/even FION: fome; 5 = A R1 1	w cone;	R2 anthers	cone; 4	R3	= Other_	
V gma V S	2 a little POLLEN P 1 = Non 3 TIGMA SH 1 = Capi	short RODUCT e; 3 = S HAPE: (4 tate; 2 =	2 = Narro R1 3 Ter/even FION: fome; 5 = A R1 1 See figure 8) = Clavate; 3 R1 1	w cone; with bundan B = Bi-le	R2 anthers R2 obed		R3 R3		R4
V gma V	2 a little POLLEN P 1 = Non 3 TIGMA SH 1 = Capi	short RODUCT e; 3 = S HAPE: (4 tate; 2 =	2 = Narro R1	w cone; with bundan B = Bi-le	R2 anthers R2 obed R2		R3 R3		R4
V gma V S (()	1 = Brown 2 2 a little POLLEN P) 1 = None 3 3 TIGMA SH 1 = Capital TIGMA Conclet the app. 46A	AAPE: (Atate; 2 =	2 = Narro R1	w cone; with bundan B = Bi-le UE: Ro	R2 anthers R2 obed R2 oyal Horticulture	Society	R3 R3 R3 R3		R4 Color Chart

TUBER CHARACTI				
PREDOMINANT; $1 = White: 2 =$		ellow; $4 = Buff$; $5 = Ta$	n: 6 - Brown, 7 - Dint	9 - D.J.
9 = Purplish-re	ed; $10 = \text{Purple}$; $11 =$	Dark purple-black; 12 =	of ther	; 8 = Red;
V 4	R1 5	TO TO	Tra Tra	
4	KI)	R2	R3	R4
PREDOMINANT S (Circle the appropri	SKIN COLOR CHART ate color chart)	VALUE: Royal Horticul	ture Society Color Chart	or Munsell Color Char
v 199C	R1 199B			1
V 1990	R1 199B	R2	R3	R4
SECONDARY SKI	N COLOR:			
1 = Absent; 2 =	= Present, please descri	be: eyes are blu	e in color	
v 2 viol	$ e _{\mathbf{R}1} _{1}$	R2	R3	R4
SECONDARY SKIP	V COLOD CHAPT VA	LUE David Hard M.		F
(Circle the appropria	te color)	LUE: Royal Horticulture	Society Color Chargor N	Iunsell Color Chart
				•
97.	3. 00-	7		
V 87A SECONDARY SKIN 1 = Eyes; 2 = E	R1 Absent PEY NEW \ N COLOR DISTRIBUT Eyebrows; 3 = Splashed	02.22.0] ION: LMC 8-0	R3 -3-2007	7 = Other
SECONDARY SKIN	PEY NEW \ N	6 provided	3-2007	
SECONDARY SKIN 1 = Eyes; 2 = E	ver new \ n N COLOR DISTRIBUT Eyebrows; 3 = Splashed	F Provided 02. 22. 07 ION: LMC 8-0 d; 4 = Scattered; 5 = S1	pectacled; 6 = Stippled;	7 = Other
SECONDARY SKIN 1 = Eyes; 2 = E V 1 SKIN TEXTURE:	per new \ no N COLOR DISTRIBUT. Eyebrows; 3 = Splashed	Provided 02-22-07 ION: LMC 8-0 d; 4 = Scattered; 5 = Sp	23-2007 pectacled; 6 = Stippled;	7 = Other
SECONDARY SKIN 1 = Eyes; 2 = E V 1 SKIN TEXTURE:	per new \ no N COLOR DISTRIBUT. Eyebrows; 3 = Splashed	F Provided 02. 22. 07 ION: LMC 8-0 d; 4 = Scattered; 5 = S1	23-2007 pectacled; 6 = Stippled;	7 = Other
SECONDARY SKIN 1 = Eyes; 2 = E V 1 SKIN TEXTURE:	per new \ no N COLOR DISTRIBUT. Eyebrows; 3 = Splashed	Provided 02-22-07 ION: LMC 8-0 d; 4 = Scattered; 5 = Sp	pectacled; 6 = Stippled; R3 = Heavily russetted; 6 =	7 = Other R4 Other
SECONDARY SKIN 1 = Eyes; 2 = E V 1 SKIN TEXTURE: 1 = Smooth; 2 =	per new \ n N COLOR DISTRIBUT Eyebrows; 3 = Splashed R1 R1 Rough (flaky); 3 = N	Provided 02-22-07 ION: LMC 8-0 d; 4 = Scattered; 5 = Sp R2 Setted; 4 = Russetted; 5	23-2007 pectacled; 6 = Stippled;	7 = Other
SECONDARY SKIN 1 = Eyes; 2 = E V 1 SKIN TEXTURE: 1 = Smooth; 2 = V 2	Per New \ No COLOR DISTRIBUT Eyebrows; 3 = Splashed R1	Provided 02-22-07 ION: LMC 8-0 d; 4 = Scattered; 5 = Sp R2 Setted; 4 = Russetted; 5	pectacled; 6 = Stippled; R3 = Heavily russetted; 6 =	7 = Other R4 Other
SECONDARY SKIN 1 = Eyes; 2 = E V 1 SKIN TEXTURE: 1 = Smooth; 2 = V 2 TUBER SHAPE: (See	Per New \ No No COLOR DISTRIBUT Eyebrows; 3 = Splashed R1	Provided 02-22-07 ION: LMC 8-0 d; 4 = Scattered; 5 = Sp R2 Setted; 4 = Russetted; 5	Page 1200 7 pectacled; 6 = Stippled; R3 = Heavily russetted; 6 = R3	7 = Other R4 Other
SECONDARY SKIN 1 = Eyes; 2 = E V	Per New \ N N COLOR DISTRIBUT Eyebrows; 3 = Splashed R1 R1 R1 3 R1 2 = Rough (flaky); 3 = N R1 2 = Round; 3 = Oval	Provided	P3-2007 pectacled; 6 = Stippled; R3 = Heavily russetted; 6 = R3 g; 6 = Other	7 = Other R4 Other R4
SECONDARY SKIN 1 = Eyes; 2 = E V 1 SKIN TEXTURE: 1 = Smooth; 2 = V 2 TUBER SHAPE: (See	Per New \ No No COLOR DISTRIBUT Eyebrows; 3 = Splashed R1	R2 R2 R2 R2	Page 1200 7 pectacled; 6 = Stippled; R3 = Heavily russetted; 6 = R3	7 = Other R4 Other
SECONDARY SKIN 1 = Eyes; 2 = E V	Per New \ N N COLOR DISTRIBUT Eyebrows; 3 = Splashed R1 R1 R1 3 R1 2 = Rough (flaky); 3 = N R1 2 = Round; 3 = Oval	Provided	P3-2007 pectacled; 6 = Stippled; R3 = Heavily russetted; 6 = R3 g; 6 = Other	7 = Other R4 Other R4
SECONDARY SKIN 1 = Eyes; 2 = E V 1 SKIN TEXTURE: 1 = Smooth; 2 = V 2 TUBER SHAPE: (Se 1 = Compressed; V 3	R1 3 Rough (flaky); 3 = N R1 3 Re figure 10) 2 = Round; 3 = Oval	Provided	P3-2007 pectacled; 6 = Stippled; R3 = Heavily russetted; 6 = R3 g; 6 = Other	7 = Other R4 Other R4
SECONDARY SKIN 1 = Eyes; 2 = E V 1 SKIN TEXTURE: 1 = Smooth; 2 = V 2 TUBER SHAPE: (Set 1 = Compressed; V 3	R1 3 Rough (flaky); 3 = N R1 3 Re figure 10) 2 = Round; 3 = Oval R1 3	Provided	P3-2007 pectacled; 6 = Stippled; R3 = Heavily russetted; 6 = R3 g; 6 = Other	7 = Other R4 Other R4
SECONDARY SKIN 1 = Eyes; 2 = E V 1 SKIN TEXTURE: 1 = Smooth; 2 = V 2 TUBER SHAPE: (Se 1 = Compressed; V 3 TUBER THICKNESS 1 = Round; 2 = 1	R1 Rough (flaky); 3 = N R1 3 Refigure 10) 2 = Round; 3 = Oval R1 3	Provided	P3-2007 pectacled; 6 = Stippled; R3 = Heavily russetted; 6 = R3 g; 6 = Other	7 = Other R4 Other R4
SECONDARY SKIN 1 = Eyes; 2 = E V 1 SKIN TEXTURE: 1 = Smooth; 2 = V 2 TUBER SHAPE: (Set 1 = Compressed; V 3	R1 3 Rough (flaky); 3 = N R1 3 Re figure 10) 2 = Round; 3 = Oval R1 3	Provided	P3-2007 pectacled; 6 = Stippled; R3 = Heavily russetted; 6 = R3 g; 6 = Other	7 = Other R4 Other R4

STD-476 (01-96)

Page 8 of 19

AVERAGE:			
V 78.68 R1 67.14	4 R2	R3	R4
RANGE:			
V 60 to 108 R1 51 to 88	8 R2 to	R3 to	R4 to
STANDARD DEVIATION:		· 	
v 10.57 R1 8.63	R2	R3	R4
AVERAGE WEIGHT OF SAMPLE TA	AKEN:		
V R1	R2	R3	R4
TUBER WIDTH (mm):			·
AVERAGE:	1 (
V 61.44 R1 65.72	R2	R3	R4
RANGE:			
V 49 to 81 R1 51 to 84	R2 to	R3 to	R4 to
STANDARD DEVIATION:			
v - 7.28 R1 5.96	R2	R3	R4
AVERAGE WEIGHT OF SAMPLE TA	KEN:		
V R1	R2	R3	R4
TUBER THICKNESS (mm):			
TUBER THICKNESS (mm): AVERAGE:			
	R2	R3	R4
AVERAGE: V 48.75 RANGE:	R2	R3	R4
AVERAGE: V 48.75 R1 51.53	R2 to	R3 to	R4 to
AVERAGE: V 48.75 RANGE:		7	
AVERAGE: V 48.75 R1 51.53 RANGE: V 36 to 61 R1 38 to 63		7	
AVERAGE: V 48.75 R1 51.53 RANGE: V 36 to 61 R1 38 to 63 STANDARD DEVIATION:	R2 to	R3 to	R4 to
AVERAGE: V 48.75 R1 51.53 RANGE: V 36 to 61 R1 38 to 63 STANDARD DEVIATION: V 4.55 R1 4.46	R2 to	R3 to	R4 to
AVERAGE: V 48.75 R1 51.53 RANGE: V 36 to 61 R1 38 to 63 STANDARD DEVIATION: V 4.55 R1 4.46 AVERAGE WEIGHT OF SAMPLE TAP	R2 to R2 KEN:	R3 to	R4 to
AVERAGE: V 48.75 R1 51.53 RANGE: V 36 to 61 R1 38 to 63 STANDARD DEVIATION: V 4.55 R1 4.46 AVERAGE WEIGHT OF SAMPLE TAP V R1 TUBER EYE DEPTH;	R2 to R2 KEN:	R3 to	R4 to

TUBER LATERAL EYE 1 = Protruding; 2 =		nediate; 4 = Deep; 5 =	Very deep	
v=occasional protru	R1 3	R2	R3	R4
NUMBER EYE / TUBER	:			
AVERAGE:				
V 11.25	R1 11.6	R2	R3	R4
RANGE:	,			
V 10 to 13 R1	10 to 13	R2 to	R3 to	R4 to
DISTRIBUTION OF TUB 1 = Predominantly api		ributed		
v 1	R1 1	R2	R3	R4
PROMINENCE OF TUBE 1 = Not prominent; 2	R EYEBROWS: = Slight prominence;	3 = Medium prominen	ce; 4 = Very prominent	; 5 = Other
v 2/3	R1 2	R2	R3	R4
* PRIMARY TUBER FLI (Circle the appropriate color	chart)	F VALUE: Royal Hortic	culture Society Color Cha	urt or Munsell Color Chart
1600	158A	R2	R3	R4
CECONDARY OF DEPART		LMG 8-03.	22.01	
SECONDARY TUBER FLI 1 = Absent; 2 = Presen				
v	R1	R2	R3	R4
SECONDARY TUBER FLE (Circle the appropriate color	ESH COLOR CHART chart)	TVALUE: Royal Hortic	ulture Society Color Cha	rt or Munsell Color Chart
V	1	R2	R3	R4
NUMBER OF TUBERS / PI 1 = Low (<8); 2 = Med		gh (>15)		
v 2	R1 2	R2	R3	R4

STD-476 (01-96)

Page 10 of 19

DISEASES REACT	ION: 0 = NOT TEST	ED; 1 = RESISTANT; 3	160000	
	5 = MODERAT	ELY SUSCEPTIBLE; 7=	s = moderately res Susceptible; 9=Hig	ISTANT; HLY SUSCEPTELE
BACTERIAL RIN	G ROT, FOLIAR REA			
v	R1	R2	R3	D4
\			IX3	R4
	G ROT, TUBER REAC			
V	R1	R2	R3	R4
LATE BLIGHT:		·		_
v 4	R1 O	R2	R3	R4
PLRV (LEAF RO)	LL):		, b	<u> </u>
V	R1	R2	R3	24
			KJ	R4
PVX:	Dt			· · · · · · · · · · · · · · · · · · ·
V	R1	R2	R3	R4
PVY:		· ·		
V	R1	R2	R3	R4
OTHER: SOFT	vot (Erwinia)			
v 4	R1 O	R2	R3	R4
OTHER:	per new infor	mation 02-22.	07	
v	R1	R2	R3	D4
PECING CHAP A COMME	<u> </u>		1.0	R4
PESTS CHARACTERIS	:			
PEST REACTION: () = NOT TESTED; 1 = 5 = MODERATELY ST	RESISTANT; 3 = MO SCEPTIBLE; 7=SUSCE	DERATELY RESISTAN	Γ;
GOLDEN NEMATO	:	umed susceptible	3	
V 7	R1 7			
	KI /	R2	R3	R4
OTHER:	:	:	<u> </u>	·
V	R1	R2	R3	R4
GENE TRAITS:	:			
INSERTION OF GEN	ES:	YES	T wa	· · · · · · · · · · · · · · · · · · ·
	<u> </u>	; 	NO	
u res, describe the g	ene(s) introduced or atta	ch information:	•	
			i	
				,

CHIEF MARKET: Processing for	r potato chip pro	oduction		
SPECIFIC GRAVIT $1 < 1.060; 2 = 1$	Y (wt. air /wt. air - wt060-1.069; 3 = 1.070-1	water) 1.079; 4 = 1.080-1.089;	5 > 1.090	
				<u> </u>
V 4	R1 4	R2	R3	R4
TOTAL GLYCOALI	KALOID CONTENT (m	g. / 100 g. fresh tuber)	· · · · · · · · · · · · · · · · · · ·	
				· [
v 5.94	R1 18.97	R2	R3	R4
OTHER QUALITY C	HARACTERISTICS: I	Describe any other quality	v characteristics that ma	ıv aid in identification
(e.g. chip-processing, t	french fry processing, ba	iking, boiling, after-cooki	ng darkening). Please a	ttach data and
corresponding protoco	01,		٠	
		·. <u></u>		
				4
•				
HEMICAL IDENTIFI Describe chemical trai	ts of the candidate varie	ty that aid in its identifica	ation (e.g. protein or DN	A electrophoresis).
Describe chemical trai Please attach data and	ts of the candidate varie the corresponding proto	ty that aid in its identifica	ation (e.g. protein or DN	A electrophoresis).
Describe chemical trai	ts of the candidate varie the corresponding proto	ty that aid in its identifica ocol.	ation (e.g. protein or DN	A electrophoresis).
Describe chemical trai Please attach data and	ts of the candidate varie the corresponding proto	ty that aid in its identifica	ation (e.g. protein or DN	A electrophoresis).
Describe chemical trai Please attach data and	ts of the candidate varie the corresponding proto	ty that aid in its identificancel.	ation (e.g. protein or DN	A electrophoresis).
Describe chemical trai Please attach data and See exhibit D.	ts of the candidate varied the corresponding proto	ocol.	ation (e.g. protein or DN	A electrophoresis).
Describe chemical trai Please attach data and See exhibit D.	ts of the candidate varie the corresponding proto	ocol.	ation (e.g. protein or DN	A electrophoresis).
Describe chemical trai Please attach data and See exhibit D.	ts of the candidate variethe corresponding proto	ocol.		A electrophoresis).
Describe chemical trai Please attach data and See exhibit D.	ts of the candidate variethe corresponding proto	eristics:		A electrophoresis).
Describe chemical trai Please attach data and See exhibit D.	ts of the candidate variethe corresponding proto	eristics:		A electrophoresis).
Describe chemical trai Please attach data and See exhibit D.	ts of the candidate variethe corresponding proto	eristics:		A electrophoresis).
Describe chemical trai Please attach data and See exhibit D.	ts of the candidate variethe corresponding proto	eristics:		A electrophoresis).
Describe chemical trai Please attach data and See exhibit D.	ts of the candidate variethe corresponding proto	eristics:		A electrophoresis).
Describe chemical trai Please attach data and See exhibit D.	ts of the candidate variethe corresponding proto	eristics:		A electrophoresis).
Describe chemical trai Please attach data and See exhibit D.	ts of the candidate variethe corresponding proto	eristics:		A electrophoresis).
Describe chemical trai Please attach data and See exhibit D.	ts of the candidate variethe corresponding proto	eristics:		A electrophoresis).
Describe chemical trai Please attach data and See exhibit D.	ts of the candidate variethe corresponding proto	eristics:		A electrophoresis).
Describe chemical trai Please attach data and See exhibit D.	ts of the candidate variethe corresponding proto	RISTICS: be useful in distinguishing		A electrophoresis).
Describe chemical trai Please attach data and See exhibit D.	ts of the candidate variethe corresponding proto	eristics:		A electrophoresis).
Describe chemical trai Please attach data and See exhibit D.	ts of the candidate variethe corresponding proto	RISTICS: be useful in distinguishing		A electrophoresis).
Describe chemical trai Please attach data and See exhibit D.	ts of the candidate variethe corresponding proto	RISTICS: be useful in distinguishing		A electrophoresis).

Exhibit C (Potato) NAME OF APPLICANT (S) VARIETY NAME TEMPORARY OR EXPERIMENTAL DESIGNATION FOR OFFICIAL USE ONLY ADDRESS (Street and No. or RD No., City, State, Zip Code, and Country) PVPO NUMBER #200200127 REFERENCE VARIETIES: Enter the reference variety name in the appropriate box. Reference Variety 4 (R4) Reference Variety 3 (R3) Application Variety (V) Reference Variety 1 (R1) Reference Variety 2 (R2) Snowden FL 2006 PLEASE READ ALL INSTRUCTIONS CAREFULLY: 1. MARKET CHARACTERISTICS: *MARKET CLASS: 1 = Yellow-flesh Tablestock 2 = Round-white Tablestock 3 = Chip-processing 4 = Frozen-processing 5 = Russet Tablestock 6 = Other R1 R2 R3 R4 2. LIGHT SPROUT CHARACTERISTICS: (See Figure 1) *LIGHT SPROUT: GENERAL SHAPE 6 = Other 1 = Spherical 4 = Broad cylindrica 5 = Narrow cylindrical 2 = Ovoid 3 = Сопіса **R2 R3** R1**R4** *LIGHT SPROUT BASE: PUBESCENCE OF TIP 2 = Weak 3 = Medium 4 = Strong 5 = Very Strong 1 = Absent R2R3**R4** *LIGHT SPROUT BASE: ANTHOCYANIN COLORATION 2 = Red-violet 4 = Other(describe) 1 = Green 3 = Blue-violet R1R2 R3 R4 *LIGHT SPROUT BASE: INTENSITY OF ANTHOCYANIN COLORATION (IF PRESENT) 1 = Absent 2 = Weak 3 = Medium 4 = Strong 5 = Very Strong R₂ R3 R4 R1* LIGHT SPROUT TIP: HABIT 2 = Intermediate 3 = Open 1 = Closed R₂ **R3 R4**

•			#	200200	0127	Exhibit C (Pot
GHT SPROUT CHARA	CTERISTICS: (continued)					
	IP: PUBESCENCE = Weak 3 = Medium	4 = Strong 5 = Very	Strong			
V	R1 3	R2	23	R4		
	TIP ANTHOCYANIN COLO = Red-violet 3 = Blue-		e)			
v 3	R1	R2	23	R4		
	P: INTENSITY OF ANTHO	OCANIN COLORATION (IF P 4 = Strong 5 = Very				
v 4	R1	R2	.3	R4		
	OOT INITIALS: FREQUEN Medium 3 = Long	NCY				
v 2	R1 2	R2	13	R4		
V 2		R2 R	23	R4		
	CS: (See Figure 2)		Spreading	R4		
ANT CHARACTERIST	CS: (See Figure 2)	ct (30-45° with ground) 7 =		R4		,
ANT CHARACTERIST	cs: (See Figure 2) ground) 5 = Semi-ere	ct (30-45° with ground) 7 =	Spreading	R4		,
ANT CHARACTERIST	CS: (See Figure 2) ground) 5 = Semi-ere	ct (30-45° with ground) 7 = R2 R	Spreading			,
ANT CHARACTERIST GROWTH HABIT: 3 = Erect (>45° with V TYPE: 1 = Stem (foliage of	cs: (See Figure 2) 1 ground) 5 = Semi-ere R1 Den, stems clearly visible)	ct (30-45° with ground) 7 = R2 R2 R2 R2 R2 R2 R2	Spreading 23 eaf (Foliage closed	R4		,
ANT CHARACTERIST GROWTH HABIT: 3 = Erect (>45° with V TYPE: 1 = Stem (foliage of	cs: (See Figure 2) 1 ground) 5 = Semi-ere R1 Den, stems clearly visible)	ct (30-45° with ground) 7 = R2 R2 R2 R2 R2 R2 R2	Spreading 23 eaf (Foliage closed	R4		,
ANT CHARACTERISTIC GROWTH HABIT: 3 = Erect (>45° with V TYPE: 1 = Stem (foliage of W MATURITY: Days	(See Figure 2) 1 ground) 5 = Semi-ere R1 Deen, stems clearly visible) R1 after planting (DAP) at vin	ct (30-45° with ground) 7 = R2 R2 R2 R2 R2 R	Spreading 23 eaf (Foliage closed	R4 d, stems hardly visible) R4		,

MATURITY CLASS:

1 = Very Early (<100 DAP) 2 = Early (100-110 DAP) 3 = Mid-season (111-120 DAP) 4 = Late (121-130 DAP) 5 = Very Late (>130 DAP).

R2



R1

R1

R2

R3


R4


R3


R4

Page 4 of 19

Figure 1: Growth Habit








7 = Spreading <30° with ground

Figure 2: Leaf Silhouette








Figure 3: Terminal Leaflet Shape / Primary Leaflet Shape



1=Narrowly Ovate



2=Medium Ovate



3=Broadly Ovate



4 = Lanceolate



5=Elliptical



6=Obovate



7=Oblong

Figure 4: Terminal Leaflet Shape of Tip / Primary Leaflet Shape of Tip




Figure 5: Terminal Leaflet Shape of Base / Primary Leaflet Shape of Base





Figure 6: Corolla Shape





stellate | > b




semi-stellate | = b



pentagonal l < b



rotate | << b



very rotate

Figure 7: Anther Shape




Figure 8: Stigma Shape




Figure 9: Distribution of Secondary Tuber Color




Figure 10: Tuber Shape












Figure 11: Leaf Dissection




Figure: 12 Stem Wings




Figure 13: Leaf Stipules:




EXHIBIT D. Additional Description of the Variety

- 1) Isozyme fingerprint of FL 2006, with reference to methodology. Comparison of fingerprint with that of Snowden, showing distinct patterns for each variety (Exhibit D-1).

 Dimension summary of 100-tuber sample of tuber dimensions of FL 2006 compared to 100 tubers
- 2) of Snowden.

EXHIBIT D-1

	Isozyme electr	\mathcal{L}	phoresis fingerprints of FL 2006 compared to Snowden.	ingerpri	nts of FL	2006 co	mpared	to Snow	den.					
Variety	Mdh-1	Mdh-1 Mdh-2	Pgdh-3 ldh-1 Pgi-1 Aps-1 Got-1 Got-2 Pgm-1 Pgm-2 Dia-1 Prx-1 Prx-3 Adh-1	ldh-1	Pgi-1	Aps-1	Got-1	Got-2	Pgm-1	Pgm-2	Dia-1	Prx-1	Prx-3	Adh-1
Snowden	1224	2222	2222	1112 2222	2222	ŀ	3344	3555	1122 2	2223 1111	1111	1	ı	ŀ
FL 2006	2222	2222	1112.	1112 1112 2224	2224	;	3344	3335	3344 3335 1111 2223	2223	i	ł	1144	ł

Source of Data: Dr. David Douches, Michigan State University 2001

Procedures and allelic designations used are according to Douches, D.S. and K. Ludlam. 1991. Electrophoretic Characterization of North American Potato Cultivars. Am. Potato J. 68:767-780.

Plant Variety Protection

FL 2006 - these tubers were grown in the Meredith field (Starks)

Date: 23 October 01

Tuber Measurements

ruper measur		
Length (mm)	Width (mm)	Depth (mm)
86	57	43
95	68	53
75	59	49
98	80	56
91	72	50
69	60	45
81	57	50
83	64	52
80	57	46
73	49	41
67	52	44
85	65	50
70	60	48
95	74	52
90	68	50
78	67	54
83	56	48
98	72	57
75	60	48
76	60	50
80	55	51
94	66	53
66	50	44
74	57	43
92	64	44
86	67	52
84	69	46
98	71	55
66	52	52
66	58	49
66	57	52
70	52	44
90	68	53
76	67	56
90	72	55
73	51	38
69	53	44
94	55	47
100	71	55
77	61	46
91	64	52
76	66	48
92	71	54

Length:			
Mean	78.68	Max	108
STD	10.57	Min	60
Width:			
Mean	61.44	Max	81
STD	7.28	Min	49
Depth:			
Mean	48.75	Max	61
STD	4.55	Min	36

#200200127

73	57	48
76	54	49
80	59	48
73	60	47
91	72	54
99	79	60
77	56	44
67	65	61
66	53	43
87	81	52
84	68	52
80	71	52
77	66	49
84	67	54
93	68	54
73	57	44
70	52	50
69	52	43
83	65	53
76	61	48
75	61	47
64	56	45
87	66	53
92	66	56
76	59	50
74	65	46
84	67	48
75	56	46
84	54	48
69	58	49
73	62	50
75	61	50
68	58	49
72	66	53
60	49	42
69	53	44
108	64	53
91	71	49
66	54	48
64	54	43
78	66	47
73	54	43
73	61	44
78	58	45
69	57	45
64	56	41
80	56	49
61	55	36
73	50	45

#200200127

62	55	43
73	58	47
63	53	49
69	58	46
69	62	46
95	72	52
83	66	52
83	70	52

Plant Variety Protection

Snowden - these tubers were grown in the Meredith field (Starks)

Date: 23 October 01

Tuber Measurements

i ubei ineasu			
			C=Compressed
88	65	57	
70	70	50	
85	69	58	
85	78	58	
85	83	57	
70	74	54	
68	74	55	C
86	74	57	
55	63	46	O
76	63	47	
56	65	50	C
71	70	51	
68	58	44	
63	63	42	
78	74	52	
73	67	51	
60	55	45	
73	73	53	
59	60	48	С
70	66	49	
66	64	49	
67	69	58	С
63	61	51	******
61	63	52	С
76	68	44	
66	56	42	
63	63	51	
65	63	44	
66	69	52	
67	67	53	
72	66	53	
64	64	49	
72	70	48	
66	59	50	
58	51	50	
61	66	59	С
80	70	54	
67	65	50	
86	78	59	
70	66	38	
73	73	58	
59	64	49	
60	63	54	С

Length:			
Mean	67.14	Max	88
STD	8.63	Min	51
· Width:	•		
Mean	65.72	Max	84
STD	5.96	Min	51
Depth:			
Mean	51.53	Max	63
STD	4.46	Min	38

67	67	51	
71	.67	53	
83	75	53	<u>'</u>
66	65	56	
72	59	44	
78	78	54	
81	74	53	
73	72	53	
71	67	51	
55	59	47	С
73	73	56	
66	69	51	C
63	70	54	С
59	59	49	
63	63	53	
64	67	51	С
66	66	53	
63	65	47	С
70	71	55	C C
55	59	51	С
63	63	46	
64	67	51	С
59	58	54	
59	59	50	
52	64	50	С
62	63	50	0 0
79	84	59	С
79	66	54	
51	59	58	
71	63	45	
70	65	56	
73	67	55	
59	61	45	С
61	59	45	
86	74	55	
59	61	58	C C
57	69	46	С
66	64	53	
60	58	49	
59	59	53	
57	58	55	С
58	61	50	C
67	67	56	
83	70	63	
59	60	48	С
60	65	54	C C
71	65	46	
60	59	48	
64	64	52	

60	64	51	С
58	64 59	54	
61	64	50	С
53	58	51	
66 66	71	56	С
66	63	51	
70	66 68	55	
66	68	55 55	

REPRODUCE LOCALLY: Include form number and edition date	on all reproductions. FORM APPROVED - OMB No. 0581-0055			
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHI	Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). The information is held confidential until the certificate is issued (7 U.S.C. 2426).			
1. NAME OF APPLICANT(S)	2 TEMPORARY DESIGNATION 3. VARIETY NAME OR EXPERIMENTAL NUMBER			
Frito-Lay North America, Inc.	1995 191.2 FL2006			
4. ADDRESS (Street and No. of R.F.D. No., City, State, and ZiP, and Country	5. TELEPHONE (include area code) 6. FAX (include area code)			
7701 Legacy Drive	(972)334-3822 (972)334-5965			
Plano, Texas 75024	#200200127			
9. Is the applicant (individual or company) a U.S. National or a	U.S. based company? If no, give name of country X YES NO			
10. Is the applicant the original owner? X YES	NO If no, please answer <u>one</u> of the following:			
a. If the original rights to variety were owned by individual	(s), is (are) the original owner(s) a U.S. National(s)?			
YES	NO If no, give name of country			
b. If the original rights to variety were owned by a compar	ny(ies), is (are) the original owner(s) a U.S. based company?			
YES NO If no, give name of country				
11. Additional explanation on ownership (If needed, use the re	verse for extra space):			
	, all rights to inventions and discoveries made Frito-Lay, Inc. are assigned to Recot, Inc., with			
PLEASE NOTE:				
Plant variety protection can only be afforded to the owners (no	t licensees) who meet the following criteria:			
 If the rights to the variety are owned by the original breeder, national of a country which affords similar protection to natio 	that person must be a U.S. national, national of a UPOV member country, or onals of the U.S. for the same genus and species.			
If the rights to the variety are owned by the company which enationals of a UPOV member country, or owned by nationals genus and species.	employed the original breeder(s), the company must be U.S. based, owned by s of a country which affords similar protection to nationals of the U.S. for the same			
3. If the applicant is an owner who is not the original owner, bo	th the original owner and the applicant must meet one of the above criteria.			
The original breeder/owner may be the individual or company vact for definitions.	who directed the final breeding. See Section 41(a)(2) of the Plant Variety Protection			
According to the Paperwork Reduction Act of 1995, an agency may not conduct or s control number. The valid OMB control number for this information collection is 058	sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB 81-0055. The time required to complete this information collection is estimated to average 6 minutes per			

response, including the time for reviewing the instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

State of Delaware Secretary of State Division of Corporations Delivered 05:24 PM 01/15/2004 FILED 05:24 PM 01/15/2004 SRV 040033329 - 2202650 FILE

STATE OF DELAWARE CERTIFICATE OF AMENDMENT OF CERTIFICATE OF INCORPORATION

Recot, Inc., a corporation organized and existing under and by virtue of the General Corporation Law of the State of Delaware.

DOES HEREBY CERTIFY:

FIRST: The Board of Directors of Recot, Inc. has duly adopted a resolution senting forth a proposed amendment of the Certificate of Incorporation of said corporation, declaring said amendment to be advisable. The resolution setting forth the proposed amendment is as follows:

RESOLVED, that the Certificate of Incorporation of this corporation be amended by changing the Article thereof numbered "ARTICLE ONE" so that, as amended, said Article shall be and read as follows:

The name of the corporation is FRITO-LAY NORTH AMERICA, INC.

SECOND: That said amendment was duly adopted in accordance with the provisions of Section 242 of the General Corporation Law of the State of Delaware.

THIRD: That the capital of said corporation shall not be reduced under or by reason of said amendment.

FOURTH: That this name change shall be effective December 29, 2003, for accounting purposes only.

FIFTH: That the Principal Administrative Office for Frito-Lay North America, Inc. shall be located at 7701 Legacy Drive, Plano, Texas 75024.

IN WITNESS WHEREOF, said Recot, Inc. has caused this certificate to be signed by Kelly Mahon Tullier, an Authorized Officer, this 20th day of January, 2004.

Authorized Officer

Title: Vice President and Treasurer Name: Kelly Mahon Tullier

EXHIBIT E. Statement of the Applicant's Ownership

Breeders employed by Frito-Lay, Inc developed the variety FL 2006, for which Plant Variety Protection is hereby sought. By agreement between Frito-Lay and its employees, all rights to inventions and discoveries made by the employees while employed by Frito-Lay, Inc. are assigned to Recot, Inc., with no ownership rights of any kind retained by employees.

REPRODUCE LOCALLY. Include form number and date on all reproductions.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is o581-0055. The time required to complete this information collection is estimated to average 5 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY PLANT VARIETY PROTECTION OFFICE BELTSVILLE, MD 20705

EXHIBIT F

	DECLARATION REGARDING DEPOSIT	
NAME OF OWNER (S)	ADDRESS (Street and No. or RD No., City, State, and Zip Code and Country)	TEMPORARY OR EXPERIMENTAL DESIGNATION
Frito Lay North America, Inc.	7701 Legacy Drive	1995 191.2
	Plano, TX 75024	VARIETY NAME FL 2006
NAME OF OWNER REPRESENTATIVE (S)	ADDRESS (Street and No. or RD No., City, State, and Zip Code and Country)	FOR OFFICIAL USE ONLY
Robert W. Hoopes	7701 Legacy Drive Plano, TX 75024	PV# 1218 0 0 2 0 0 1 2 7

I do hereby declare that during the life of the certificate a viable sample of propagating material of the subject variety will be deposited, and replenished as needed periodically, in a public repository in the United States in accordance with the regulations established by the Plant Variety Protection Office.